

BOEHRINGER INGELHEIM VETMEDICA, INC.

2621 NORTH BELT HIGHWAY, ST. JOSEPH, MO, 64506-2002

Telephone: 800-325-9167

Fax: 816-236-2717

Website: www.bevaccinesmart.com

Website: www.bi-vetmedica.com

Website: www.haveweseenyourcatlatelately.com

Website: www.metacam.com

Website: www.mycathasdiabetes.com

Website: www.prrsresearch.com

Website: www.prozinc.us

Website: www.vetera-vaccines.com

Website: www.vetmedin-us.com

Website: www.yourdogsheart.com

Email: info@productionvalues.us



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

Diaque®

Boehringer Ingelheim

Nutritional supplement for calves and other species

Source of energy and electrolytes

Compatible with milk, milk replacer or water

Diaque has a high energy content

Diaque is a unique formula containing a patented hydrophobic citrus fiber.

Diaque supports fast absorption of electrolytes and water. Its unique three step buffering system allows it to be fed with milk, milk replacer or water. Diaque may be used during times of stress due to changes in feed or environment. Diaque can be mixed in the normal milk feeding.

Diaque Feeding Directions:

A 50 g scoop is enclosed for measuring Diaque (6.6 lb, 22 lb sizes).

Calves: For oral use only. Mix 1 packet or 2 scoops (100 g) of Diaque in 2 quarts (1891 ml) of warm water, milk or milk replacer. Feed immediately. For larger calves or those requiring more energy, mix 1 1/2 packets or 3 scoops (150 g) Diaque in 3 quarts (2837 ml) of warm water, milk or milk replacer. Feed every 8 to 10 hours for 24 to 48 hours. Discard any unconsumed mixture each day.

Pigs: Mix 1 packet or 2 scoops (100 g) of Diaque in 2 quarts (1891 ml) of warm water for a litter of 8 pigs. Feed in 4 feedings per day from a pan. Discard any remaining portion at each feeding. Normal feeding is for 2 days.

Foals: Mix 1 packet or 2 scoops (100 g) of Diaque in 2 quarts (1891 ml) of warm water. Feed this amount to foal twice daily. May be fed with bucket or bottle. Amount recommended is for a 110 lb foal. Adjust for larger or smaller animals. Normal feedings are for 2 days.

Lambs, Kids, Crias: Remove from dam. Each feeding for a 10 lb animal should consist of 1 tablespoon (10 g) of Diaque mixed in 1/2 pint (236 ml) of warm water. The mixture may be fed with bucket or bottle. Adjust amounts for larger or smaller animals. Normal feedings are for 2 days.

Caution:

For animal use only.

Keep out of reach of children.

Store in a dry place below 86° F (30° C).

Guaranteed analysis:

Crude Protein, minimum	2.2%
------------------------	------

Crude Fat, minimum	4.6%
Crude Fiber, maximum	4.5%
Salt, minimum	4.5%
Salt, maximum	5.3%
Sodium, minimum	3.4%
Sodium, maximum	5.9%
Potassium, minimum	1.0%
Ash, maximum	15.8%
Glucose, minimum	53.5%

Ingredients:

Glucose, dried citrus pulp, lecithin, salt, sodium bicarbonate, silicon dioxide, potassium chloride, sodium citrate, sodium acetate, dried yeast, glycine (aminoacetic acid), xanthan gum (stabilizer), natural and artificial flavors and ethoxyquin (preservative).

Product of Denmark

Manufactured by: Boehringer Ingelheim Denmark A/S, DK-2100 Copenhagen O E

Manufactured for: Boehringer Ingelheim Vetmedica, Inc., St. Joseph, MO 64506 U.S.A.

Toll Free 1-866-638-2226

Diaque is a registered trademark of Boehringer Ingelheim Vetmedica GmbH, used under license.

Net Content/Weight	
24 x 3.5 oz (100 g) packets	434201-01 4690e-1411
6.6 lb (3 kg)	434204-01 5956-1411
22 lb (10 kg)	434205-01 5599-1103

CPN: 1028159.5