

PANACUR C CANINE- fenbendazole granule
Merck Sharp & Dohme Corp.

panacur[®] c
(fenbendazole)

Drug Facts

Active ingredients (in each dosage unit):

Fenbendazole Granules 22.2% (222 mg/g)

Purpose: Dewormer for Dogs Only

Uses: For the treatment and control of Roundworms (*Toxocara canis*, *Toxascaris leonina*), Hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*), Whipworms (*Trichuris vulpis*), and Tapeworms (*Taenia pisiformis*).

Human Warning: Keep this and all medication out of the reach of children.

Adverse Reactions: In US clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with use of the product.

When using this product: Panacur[®] C is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

Directions:

The daily dose for Panacur[®] C is 50 mg/kg (22.7 mg/lb) of body weight. Please refer to the following dosing table for help in finding the right dose for your dog.

Dosing Table

Dog Weight	Packet Size *
10 pounds	1 gram
20 pounds	2 gram
30 pounds	1 gram + 2 gram
40 pounds	4 gram
50 pounds	1 gram + 4 gram
60 pounds	2 gram + 4 gram
80 pounds	Two 4 gram
Over 80 pounds	Use combinations to obtain recommended daily dose.

* **Packet size is the daily dose. The dog must be treated with this dose for 3 days in a row.**

- You should weigh your dog to make sure you are using the right size and number of packets. If your dog's weight is in-between the suggested dosing sizes, it is safe to use the next higher size. For example, a 15-pound dog should be treated with the 2-gram packet.
- Give Panacur[®] C to your dog by mixing the daily dose with a small amount of the usual food. Make sure your dog eats all of the medicated food. You may need to moisten dry dog food to aid mixing.
- Repeat the daily dose for three days in a row.

Other Information:

Diagnosis of Parasites: Specific diagnoses require laboratory testing. Consult your veterinarian for

help in the diagnosis, treatment, and control of parasitism. Dogs can become infected with several different kinds of tapeworms. Panacur[®] C only kills *Taenia* species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Panacur[®] C, consult your veterinarian.

Recommended Deworming Schedule:

Deworming schedules may vary depending on the climate where you live and the activity of your dog. The following schedule should be used as general guidance. Newly weaned pups (6 to 8 weeks of age) should be dewormed at 6, 8, 10, and 12 weeks of age. Treat the dam at the same time as the puppies. Dogs over six months of age should be dewormed at least twice each year. Each deworming requires 3 daily treatments. (See Directions)

Storage: Store at controlled room temperature 68-77°F (20-25°C).

Questions? Comments?

To report a suspected adverse reaction or to obtain product information including material safety data sheets (MSDS), call 1-800-441-8272.

Manufactured by:

Intervet GesmbH

Vienna, Austria

Made in Austria

Distributed by:

Intervet Inc (d/b/a Merck Animal Health)

Summit, NJ 07901

www.intervetusa.com

NADA # 121-473, Approved by FDA

446600-C

Net Weight: 6 grams

PRINCIPAL DISPLAY PANEL - 2 g Packet Carton

MERCK

Animal Health

panacur[®] c

(fenbendazole)

Canine

Dewormer

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Package
Contents:
Three 2-gram
packets

Each Packet
Treats 20 lbs.



panacur[®]C

(fenbendazole)

Canine Dewormer



Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Drug Facts

Active ingredients (in each dosage unit):

Fenbendazole Granules 22.2% (222 mg/g)

Purpose: Dewormer for Dogs Only

Uses: For the treatment and control of Roundworms (*Toxocara canis*, *Toxascaris leonina*), Hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*), Whipworms (*Trichuris vulpis*), and Tapeworms (*Taenia pisiformis*).

Human Warning: Keep this and all medication out of the reach of children.

Adverse Reactions: In US clinical studies 3 of 240 dogs (about 1% of the treated dogs) had vomiting associated with use of the product.

When using this product: Panacur[®] C is safe for use in puppies 6 weeks or older and adult dogs, including pregnant bitches. Do not deworm a dog or puppy that is sick. Consult your veterinarian for diagnosis of the illness.

Lot:/Exp: will be printed on the line in this area
lacquer- and colorfree

113678 P2

Directions:

The daily dose for Panacur[®] C is 50 mg/kg (22.7 mg/lb) of body weight. Please refer to the following dosing table for help in finding the right dose for your dog.

Dosing Table

Dog Weight	Packet Size*
10 pounds	1 gram
20 pounds	2 gram
30 pounds	1 gram + 2 gram
40 pounds	4 gram
50 pounds	1 gram + 4 gram
60 pounds	2 gram + 4 gram



80 pounds	Two 4 gram
Over 80 pounds	Use combinations to obtain recommended daily dose.

*Packet size is the daily dose. The dog must be treated with this dose for 3 days in a row.

- You should weigh your dog to make sure you are using the right size and number of packets. If your dog's weight is in-between the suggested dosing sizes, it is safe to use the next higher size. For example, a 15-pound dog should be treated with the 2-gram packet.
- Give Panacur® C to your dog by mixing the daily dose with a small amount of the usual food. Make sure your dog eats all of the medicated food. You may need to moisten dry dog food to aid mixing.
- Repeat the daily dose for three days in a row.

Other Information:

Diagnosis of Parasites: Specific diagnoses require laboratory testing. Consult your veterinarian for help in the diagnosis, treatment, and control of parasitism. Dogs can become infected with several different kinds of tapeworms. Panacur® C only kills *Taenia* species of tapeworms. If you continue to see tapeworm segments in your dog's stool after treatment with Panacur® C, consult your veterinarian.

Recommended Deworming Schedule:

Deworming schedules may vary depending on the climate where you live and the activity of your dog. The following schedule should be used as general guidance. Newly weaned pups (6 to 8 weeks of age) should be dewormed at 6, 8, 10, and 12 weeks of age. Treat the dam at the same time as the puppies. Dogs over six months of age should be dewormed at least twice each year. Each deworming requires 3 daily treatments. (See Directions)

Storage: Store at controlled room temperature 68-77°F (20-25°C).

Questions? Comments?

To report a suspected adverse reaction or to obtain product information including material safety data sheets (MSDS), call 1-800-441-8272.

Manufactured by:
Intervet GesmbH
Vienna, Austria
Made in Austria

Distributed by:
Intervet Inc (d/b/a Merck Animal Health)
Summit, NJ 07901
www.intervetusa.com

NADA # 121-473, Approved by FDA

446600-C

Net Weight: 6 grams



PSA020A2B_01_2
89x13x76mm

3280 black
L114B
x=10; y=15

PANACUR C CANINE

fenbendazole granule

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:57926-481
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FENBENDAZOLE (UNII: 621BVT9M36) (FENBENDAZOLE - UNII:621BVT9M36)	FENBENDAZOLE	222 mg in 1 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57926-481-01	3 in 1 CARTON		
1		2 g in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA121473	01/05/1979	

