

Label

White papers

Coupons

Other

Image

ZOETIS INC.

333 PORTAGE STREET, KALAMAZOO, MI, 49007

Telephone: 269-359-4414

Customer Service: 888-963-8471

Website: www.zoetis.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

Lincomix® Zoetis

brand of lincomycin injection

For Intramuscular Use in Swine Only

LINCOMIX Injectable contains lincomycin hydrochloride, an antibiotic produced by *Streptomyces lincolnensis* var. *lincolnensis*, which is chemically distinct from all other clinically available antibiotics and is isolated as a white crystalline solid.

INDICATIONS FOR SWINE

LINCOMIX Injectable is indicated for the treatment of infectious forms of arthritis caused by organisms sensitive to its activity. This includes most of the organisms responsible for the various infectious arthritides in swine, such as the staphylococci, streptococci, *Erysipelothrix* and *Mycoplasma* spp. It is also indicated for the treatment of mycoplasma pneumonia.

CONTRAINDICATIONS

As with all drugs, the use of LINCOMIX Injectable is contraindicated in animals previously found to be hypersensitive to the drug.

WARNING

Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment. **Not for human use.**

CAUTION

If no improvement is noted within 48 hours, consult a veterinarian.

ADVERSE REACTIONS

The intramuscular administration to swine may cause a transient diarrhea or loose stools. Although this effect has rarely been reported, one must be alert to the possibility that it may occur.

Should this occur, it is important that the necessary steps be taken to prevent the effects of dehydration.

DOSAGE AND ADMINISTRATION

For arthritis or mycoplasma pneumonia-5 mg per pound of body weight intramuscularly once daily for three to seven days as needed. When using LINCOMIX Injectable containing 100 mg/mL, 1 mL/20 lb body weight will provide 5 mg/lb. When using LINCOMIX Injectable containing 300 mg/mL, 1 mL/60 lb body weight will provide 5 mg/lb.

For optimal results, initiate treatment as soon as possible.

As with any multi-dose vial, practice aseptic techniques in withdrawing each dose. Adequately clean and disinfect the vial closure prior to entry with a sterile needle and syringe. No vial closure should be entered more than 20 times.

HOW SUPPLIED

LINCOMIX Injectable is available in two concentrations: 300 mg/mL and 100 mg/mL.

300 mg/mL: For use in swine weighing 300 pounds or more.

Each mL contains lincomycin hydrochloride equivalent to lincomycin, 300 mg; also Benzyl Alcohol, 9.45 mg added as preservative. Supplied in 100 mL vials.

100 mg/mL: Each mL contains lincomycin hydrochloride equivalent to lincomycin, 100 mg; also Benzyl Alcohol, 9.45 mg added as preservative. Supplied in 100 mL vials.

Store at controlled room temperature 20°-25°C (68°-77°F), with excursions between 15°-40°C (59°-104°F).

Use contents within 28 days of first vial broach.

Approved by FDA under NADA # 034-025

Distributed by: Zoetis Inc., Kalamazoo, MI 49007

Revised: May 2019

40027759

CPN: 3690475.3

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

Copyright 2021 - Animalytix LLC