

Strongid® C/C 2X™

Strongid® C, Strongid C 2X™ (pyrantel tartrate equine wormer)

Mixing, Use and Feeding Directions

MIXING AND FEEDING DIRECTIONS					
<p>Strongid C or Strongid C 2X (pyrantel tartrate) is to be administered on a continuous basis either as a top-dress or mixed in the horse's daily grain ration at the rate of 1.2 mg pyrantel tartrate per pound of body weight daily.</p> <p>To achieve this dose administer:</p> <ul style="list-style-type: none"> • 1 oz of Strongid C per 250 lb of body weight. (A Strongid C measuring cup is enclosed.) • 0.5 oz of Strongid C 2X per 250 lb of body weight. (A Strongid C 2X measuring cup is enclosed. Please note it is different in size and shape from the original Strongid C measuring cup due to the higher concentration of Strongid C 2X.) 					
TOP-DRESS DIRECTIONS					
Lb Body Weight	Ounces per day of Strongid C		Ounces per Day of Strongid C 2X		
250	1		0.5		
500	2		1.0		
750	3		1.5		
1000	4		2.0		
1250	5		2.5		
<p>Strongid C or Strongid C 2X should be administered for the entire period that the animal is at risk to internal parasites. Unprotected animals that have grazed may already have an established <i>S. vulgaris</i> larval infestation. Before administering Strongid C or Strongid C 2X, these animals should be treated with a therapeutic dose of a larvicidal product.</p>					
<p>Foals may be administered Strongid C or Strongid C 2X as soon as consistent intake of grain mix is occurring. This is generally between two to three months of age.</p>					
<p>Strongid C or Strongid C 2X may be used in mares at any stage of pregnancy or lactation. Stallion fertility is not affected by the use of Strongid C or Strongid C 2X.</p>					
MEDICATED GRAIN MIX DIRECTIONS					
Lb of Medicated Grain Mix per 100 lb of Body Weight	Strongid C		Strongid C 2X		Concentration Grams per Ton
	Lb of Strongid C	Lb of Non-Medicated Feed	Lb of Strongid C 2X	Lb of Non-Medicated Feed	
2.0	25	1975	12.5	1987.5	120
1.5	33	1967	16.5	1983.5	160
1.0	50	1950	25.0	1975	240
0.5	100	1900	50.0	1950	480
0.2	250	1750	125.0	1875	1200

GUARANTEED ANALYSIS		
	Strongid C	Strongid C 2X
Crude Protein, not less than	12.5%	11.5%
Crude Fat, not less than	2.0%	1.5%
Crude Fiber, not more than	22.0%	22.0%
INGREDIENTS		
Dehydrated Alfalfa Meal, Wheat Middlings, Ground Corn (Strongid C only), Cane Molasses, and Preserved with Propionic Acid.		

Warning

Not for use in horses intended for food.

Caution

Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment and control of parasitism. Do not mix in feeds containing bentonite.

Strongid® is a registered trademark of Pfizer Inc



Animal Health

