

STRONGID® PASTE

(pyrantel pamoate)

Paste

Description

STRONGID® (pyrantel pamoate) Paste is a potent equine wormer containing pyrantel, a compound from the tetrahydropyrimidine class. STRONGID Paste removes and controls most common and damaging internal parasites including bloodworms, small strongyles, pinworms, and large roundworms. STRONGID Paste is a pale-yellow-to-buff paste containing 43.9% w/w pyrantel pamoate in an inert vehicle. Each syringe contains 3.6 grams pyrantel base in 23.6 g (20 mL) paste. Each mL contains 180 mg pyrantel base as pyrantel pamoate.



Approved Uses

STRONGID Paste is approved for:

- The removal and control of mature infections of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); pinworms (*Oxyuris equi*); large roundworms (*Parascaris equorum*); and small strongyles in horses and ponies.

Key Features

- Removes and controls a variety of equine worms
- Effective against benzimidazole-resistant small strongyles
- Chemical class, unrelated to other classes of equine anthelmintics
- Use in rotational deworming programs
- Convenient disposable syringe treats 1,200 lb body weight
- Wide margin of safety

Packaging

20-mL tubes

Dosage and Administration

- STRONGID Paste is to be administered as a single oral dose of 3 mg pyrantel base per pound of body weight. The syringe has four (4) weight mark increments. Each weight mark indicates the recommended dose for 300 pounds of body weight.
- Note: Position screw-gauge over appropriate mark on plunger. Each mL contains 180 mg pyrantel base as pyrantel pamoate.
- For maximum control of parasitism, it is recommended that foals (2 to 8 months of age) be dosed every four (4) weeks. To minimize the potential source of infection that the mare may pose to the foal, the mare should be treated one (1) month prior to the anticipated foaling date followed by retreatment 10 days to two (2) weeks after birth of foal. Horses and ponies over eight (8) months of age should be routinely dosed every six (6) weeks.

- After removing the cap, the paste should be deposited on the dorsum of the tongue. Introduce the nozzle end of the syringe at the corner of the mouth. Direct the syringe backwards and depress the plunger to deposit the paste onto the tongue. Given in this manner, it is unlikely that rejection of the paste will occur. Raising the horse's head sometimes assists in the swallowing process. When only part of the paste has been used, replace the cap on the syringe nozzle.

Precautions

- Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.
- Not for use in horses intended for food.
- Keep out of the reach of children.
- It is recommended that severely debilitated animals not be treated with this preparation.